

# Immunomodulating and Revascularizing Activity of *Kalanchoe pinnata* Synergize with Fungicide Activity of Biogenic Peptide Cecropin P1

Zakharchenko N., Belous A., Biryukova Y., Medvedeva O., Belyakova A., Masgutova G.,  
Trubnikova E., Buryanov Y., Lebedeva A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2017 N. S. Zakharchenko et al. Previously transgenic *Kalanchoe pinnata* plants producing an antimicrobial peptide cecropin P1 (CecP1) have been reported. Now we report biological testing *K. pinnata* extracts containing CecP1 as a candidate drug for treatment of wounds infected with *Candida albicans*. The drug constitutes the whole juice from *K. pinnata* leaves (not ethanol extract) sterilized with nanofiltration. A microbicide activity of CecP1 against an animal fungal pathogen in vivo was demonstrated for the first time. However, a favorable therapeutic effect of the transgenic *K. pinnata* extract was attributed to a synergism between the fungicide activity of CecP1 and wound healing (antiscar), revascularizing, and immunomodulating effect of natural biologically active components of *K. pinnata*. A commercial fungicide preparation clotrimazole eliminated *C. albicans* cells within infected wounds in rats with efficiency comparable to CecP1-enriched *K. pinnata* extract. But in contrast to *K. pinnata* extract, clotrimazole did not exhibit neither wound healing activity nor remodeling of the scar matrix. Taken together, our results allow assumption that CecP1-enriched *K. pinnata* extracts should be considered as a candidate drug for treatment of dermatomycoses, wounds infected with fungi, and bedsores.

<http://dx.doi.org/10.1155/2017/3940743>

---

## References

- [1] S. El Abdellaoui, E. Destandau, A. Toribio et al., "Bioactive molecules in *Kalanchoe pinnata* leaves: extraction, purification, and identification," *Analytical and Bioanalytical Chemistry*, vol. 398, no. 3, pp. 1329-1338, 2010.
- [2] E. A. Suptel, V. F. Lapchik, N. B. Zagornaia, and V. P. Shirobokov, "Antimicrobial activity of *kalanchoe* juice relative to dysentery bacteria," *Mikrobiologicheskii Zhurnal*, vol. 42, no. 1, pp. 86-90, 1980.
- [3] M. M. Bass, A. A. Fedorovskii, L. V. Kobernichenko, E. F. Salivon, and V. Pinskii, "New drug preparation-*kalanchoe* juice," *Farmatsevtichnyi Zhurnal*, vol. 25, no. 3, pp. 89-93, 1970.
- [4] M. R. Jaeger Greer, R. G. Cates, F. B. Johnson, D. Lamnaouer, and L. Ohai, "Activity of acetone and methanol extracts from thirty-one medicinal plant species against herpes simplex virus types 1 and 2," *Pharmaceutical Biology*, vol. 48, no. 9, pp. 1031-1037, 2010.
- [5] V. P. Shirobokov, A. I. Evtushenko, V. F. Lapchik, D. N. Shirobokova, and E. A. Suptel, "Antiviral activity of representatives of the family Crassulaceae," *Antibiotiki*, vol. 26, no. 12, pp. 897-900, 1981.
- [6] F. G. Urmenyi, G. D. Saraiva, L. M. Casanova et al., "Anti-HSV-1 and HSV-2 flavonoids and a new kaempferol triglycoside from the medicinal plant *Kalanchoe daigremontiana*," *Chemistry & Biodiversity*, vol. 13, no. 12, pp. 1707-1714, 2016.

- [7] S. A. Da Silva, S. S. Costa, S. C. Mendonca, E. M. Silva, V. L. Moraes, and B. Rossi-Bergmann, "Therapeutic effect of oral *Kalanchoe pinnata* leaf extract in murine leishmaniasis," *Acta Tropica*, vol. 60, no. 3, pp. 201-210, 1995.
- [8] V. Munoz, M. Sauvain, G. Bourdy et al., "The search for natural bioactive compounds through a multidisciplinary approach in Bolivia. Part II. Antimalarial activity of some plants used by Mosetene Indians," *Journal of Ethnopharmacology*, vol. 69, no. 2, pp. 139-155, 2000.
- [9] V. Schuler, K. Suter, K. Furer et al., "Bryophyllum pinnatum inhibits detrusor contractility in porcine bladder strips-a pharmacological study towards a new treatment option of overactive bladder," *Phytomedicine*, vol. 19, no. 10, pp. 947-951, 2012.
- [10] G. Umbuzeiro-Valent, D. A. Roubicek, and E. M. Haebisch, "Mutagenic and antimutagenic evaluation of the juice of the leaves of *Bryophyllum calycinum* (*Kalanchoe pinnata*), a plant with antihistamine activity," *Environmental and Molecular Mutagenesis*, vol. 33, no. 4, pp. 325-327, 1999.
- [11] N. P. Yadav and V. K. Dixit, "Hepatoprotective activity of leaves of *Kalanchoe pinnata* Pers," *Journal of Ethnopharmacology*, vol. 86, no. 2-3, pp. 197-202, 2003.
- [12] T. Ibrahim, J. M. Cunha, K. Madi, L. M. da Fonseca, S. S. Costa, and V. L. Goncalves Koatz, "Immunomodulatory and antiinflammatory effects of *Kalanchoe brasiliensis*," *International Immunopharmacology*, vol. 2, no. 7, pp. 875-883, 2002.
- [13] R. H. Mourao, F. O. Santos, E. M. Franzotti, M. P. Moreno, and A. R. Antonioli, "Antiinflammatory activity and acute toxicity (LD50) of the juice of *Kalanchoe brasiliensis* (comb.) leaves picked before and during blooming," *Phytotherapy Research*, vol. 13, no. 4, pp. 352-354, 1999.
- [14] A. C. Ferreira, D. Rosenthal, and D. P. Carvalho, "Thyroid peroxidase inhibition by *Kalanchoe brasiliensis* aqueous extract," *Food and Chemical Toxicology*, vol. 38, no. 5, pp. 417-421, 2000.
- [15] A. I. Baginskaya and . . Leskova, "The experimental substantiation of application Kalanchin, the new medicinal drag from *Kalanchoe pinnata*," , pp. 411-414, Moscow, 2006.
- [16] U. Supratman, T. Fujita, K. Akiyama et al., "Anti-tumor promoting activity of bufadienolides from *Kalanchoe pinnata* and *K. daigremontiana* x *tubiflora*," *Bioscience, Biotechnology, and Biochemistry*, vol. 65, no. 4, pp. 947-949, 2001.
- [17] R. B. Kutsik and B. M. Zuzuk, "*Kalanchoe pinnata* (Lam.) Pers.," *Provizor*, vol. 6, no. 5, 2004.
- [18] K. Furer, M. Raith, R. Brenneisen et al., "Two new flavonol glycosides and a metabolite profile of *Bryophyllum pinnatum*, a phytotherapeutic used in obstetrics and gynaecology," *Planta Medica*, vol. 79, no. 16, pp. 1565-1571, 2013.
- [19] K. Adenike and O. B. Eretan, "Purification and partial characterization of a lectin from the fresh leaves of *Kalanchoe crenata* (Andr.) Haw," *Journal of Biochemistry and Molecular Biology*, vol. 37, no. 2, pp. 229-233, 2004.
- [20] A. I. Evtushenko, "Lectins in virology," *Voprosy Virusologii*, vol. 27, no. 2, pp. 132-137, 1982.
- [21] K. Furer, A. P. Simoes-Wust, U. von Mandach, M. Hamburger, and O. Potterat, "*Bryophyllum pinnatum* and related species used in anthroposophic medicine: constituents, pharmacological activities, and clinical efficacy," *Planta Medica*, vol. 82, no. 11-12, pp. 930-941, 2016.
- [22] M. F. Muzitano, M. C. Bergonzi, G. O. De Melo et al., "Influence of cultivation conditions, season of collection and extraction method on the content of antileishmanial flavonoids from *Kalanchoe pinnata*," *Journal of Ethnopharmacology*, vol. 133, no. 1, pp. 132-137, 2011.
- [23] B. S. Nayak, J. R. Marshall, and G. Isitor, "Wound healing potential of ethanolic extract of *Kalanchoe pinnata* Lam. leaf-a preliminary study," *Indian Journal of Experimental Biology*, vol. 48, no. 6, pp. 572-576, 2010.
- [24] V. F. Lapchik and T. V. Yurchishina, *Lectin Isolation from Kalanchoe blossfeldiana*, Kiev State University, USSR, 1986.
- [25] S. S. Costa, A. Jossang, B. Bodo, M. L. Souza, and V. L. Moraes, "Patuletin acetylramnosides from *Kalanchoe brasiliensis* as inhibitors of human lymphocyte proliferative activity," *Journal of Natural Products*, vol. 57, no. 11, pp. 1503-1510, 1994.
- [26] S. A. Da-Silva, S. S. Costa, and B. Rossi-Bergmann, "The antileishmanial effect of *Kalanchoe* is mediated by nitric oxide intermediates," *Parasitology*, vol. 118, Part 6, pp. 575-582, 1999.
- [27] N. S. Zakharchenko, A. A. Lebedeva, O. V. Furs et al., "Producing marker-free *Kalanchoe* plants expressing antimicrobial peptide cecropin P1 gene," *Russian Journal of Plant Physiology*, vol. 63, no. 2, p. 9, 2016.
- [28] R. Eckert, "Road to clinical efficacy: challenges and novel strategies for antimicrobial peptide development," *Future Microbiology*, vol. 6, no. 6, pp. 635-651, 2011.
- [29] J. Xu, F. Zhong, Y. Zhang et al., "Construction of *Bacillus subtilis* strain engineered for expression of porcine betadefensin-2/cecropin P1 fusion antimicrobial peptides and its growth-promoting effect and antimicrobial activity," *Asian-Australas Journal Animal Sciences*, vol. 30, no. 4, pp. 576-584, 2016.

- [30] A. G. Khanin, P. I. Tolstykh, L. F. Muliaev, and A. P. Levshenko, "Healing of suppurative wounds during treatment with proteolytic enzymes of bacterial origin," *Sovetskaia Meditsina*, vol. 42, no. 3, pp. 119-124, 1976.
- [31] K. Abaev lu, "The technique and technicism in surgery," *Vestnik Khirurgii Imeni I. I. Grekova*, vol. 169, no. 1, pp. 144-147, 2010.
- [32] N. Takzaree, A. Hadjiakhondi, G. Hassanzadeh, M. R. Rouini, and A. Manayi, "Synergistic effect of honey and propolis on cutaneous wound healing in rats," *Acta Medica Iranica*, vol. 54, no. 4, pp. 233-239, 2016.
- [33] B. M. Kostiuichenok, V. P. Tumanov, L. I. Muzykant, V. F. Khotinian, and A. Amiraslanov lu, "Morphological characteristics of healing of extensive purulent wounds during treatment in the controlled bacteria-free environment," *Biulleten' Eksperimental'no Biologii i Meditsiny*, vol. 94, no. 9, pp. 122-125, 1982.
- [34] A. Ebbensgaard, H. Mordhorst, M. T. Overgaard, C. G. Nielsen, F. M. Aarestrup, and E. B. Hansen, "Comparative evaluation of the antimicrobial activity of different antimicrobial peptides against a range of pathogenic bacteria," *PLoS One*, vol. 10, no. 12, article e0144611, 2015.